

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/GB05/000024

International filing date: 07 January 2005 (07.01.2005)

Document type: Certified copy of priority document

Document details: Country/Office: GB
Number: 0401988.1
Filing date: 30 January 2004 (30.01.2004)

Date of receipt at the International Bureau: 09 February 2005 (09.02.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



PCT/GB2005/000024.



INVESTOR IN PEOPLE


The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.


In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 25 January 2005



30JAN04 E869435-1 D02903
P01/7700 0:00-0401988.1 ACCOUNT CHA

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference	11344P5 GB/JCM		
2. Patent application number (The Patent Office will fill this part in)	0401988.1		30 JAN 2004
3. Full name, address and postcode of the or of each applicant (underline all surnames)	Reckitt Benckiser (Australia) Pty Limited 44 Wharf Road West Ryde New South Wales 2114 AUSTRALIA		
Patents ADP number (if you know it)	07954431001 ✓		
If the applicant is a corporate body, give the country/state of its incorporation	Australia		
4. Title of the invention	Rodenticide		
5. Name of your agent (if you have one)	John Crawford McKnight		
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	Reckitt Benckiser plc Group Patents Department Dansom Lane Hull HU8 7DS UNITED KINGDOM		
Patents ADP number (if you know it)	07799521001 ✓		
6. Priority: Complete this section if you are declaring priority from one or more earlier patent applications, filed in the last 12 months.	Country	Priority application number (if you know it)	Date of filing (day / month / year)
7. Divisionals, etc: Complete this section only if this application is a divisional application or resulted from an entitlement dispute (see note f)	Number of earlier UK application	Date of filing (day / month / year)	
8. Is a Patents Form 7/77 (Statement of inventorship and of right to grant of a patent) required in support of this request?	Yes		
Answer YES if:			
a) any applicant named in part 3 is not an inventor, or			
b) there is an inventor who is not named as an applicant, or			
c) any named applicant is a corporate body.			
Otherwise answer NO (See note d)			

Patents Form 1/77

9. Accompanying documents: A patent application must include a description of the invention. Not counting duplicates, please enter the number of pages of each item accompanying this form:

Continuation sheets of this form

Description	25
Claim(s)	5
Abstract	1
Drawing(s)	

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for a preliminary examination and search (Patents Form 9/77)

One

Request for a substantive examination (Patents Form 10/77)

One

Any other documents (please specify)

FS1

11. I/We request the grant of a patent on the basis of this application.

Signature(s)

John Crawford McKnight
John Crawford McKnight

Date 26 January 2004

12. Name, daytime telephone number and e-mail address, if any, of person to contact in the United Kingdom

John McKnight (01482) 583719
john.mcknight@reckittbenckiser.com

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered YES in part 8, a Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- Part 7 should only be completed when a divisional application is being made under section 15(4), or when an application is being made under section 8(3), 12(6) or 37(4) following an entitlement dispute. By completing part 7 you are requesting that this application takes the same filing date as an earlier UK application. If you want the new application to have the same priority date(s) as the earlier UK application, you should also complete part 6 with the priority details.

Rodenticide

Technical Field

This invention relates to rodenticidal compositions containing systemic insecticides and in particular rodenticidal compositions containing the systemic insecticide fipronil (5-amino-3-cyano-1-(2,6-dichloro-4-trifluoro-methylphenyl)-4-trifluoromethylsulfinylpyrazole).

Background Art

Rodents are often host to a range of parasitic arthropods including fleas and ticks. These can generally move from one host to another and if no host is available, they can survive for extended periods until a new host can be found. The manner in which this occurs varies from species to species.

Fleas and ticks can be vectors of organisms causing a range of diseases such as Lyme disease, plague, Rocky Mountain spotted fever, Colorado tick fever, Kyansanur Forest disease, Kerneroyo, Powassan encephalitis, Russian spring-summer encephalitis, Crimean-Congo haemorrhagic fever, tick-borne encephalitis, Mediterranean spotted fever, boutonneuse fever, Q fever, North Asian tick typhus, Queensland tick typhus, murine typhus, tick-bite fever, tularaemia, relapsing fever, ehrlichiosis and babesiosis. Tick toxins may also result in paralysis of some hosts.

At the very least, ticks and fleas cause significant irritation and discomfort to a host animal.

When rodents are killed, the vector fleas and ticks seek a new host and may transfer to domestic stock, domestic pets and humans causing discomfort and spreading disease. It is desirable therefore to control parasitic arthropods in conjunction with controlling their host rodents.

Traditionally, insecticides are dusted or sprayed just before or just after rodenticide application or an insect powder is applied in a bait box where rodents enter to reach the rodenticide. Ensuring a lethal dose of insecticide to target fleas/ticks is difficult and the possibility of killing non-target insects is high.

Disclosure of Invention

More convenient would be a combination of an insecticide and a rodenticide in a single bait, the insecticide becoming systemic (i.e. available in the blood of the host) after ingestion of the bait. It is desirable that the arthropods die before the host rodent or at least take on a lethal dose of insecticide to prevent successful transfer to a new host.

Many rodents are difficult to kill because they are naturally suspicious and will not easily take a bait. When incorporating an insecticide into a bait, the bait must remain sufficiently palatable so that the rodent will accept it and ingest a lethal dose. Accordingly, it is desirable to provide an insecticide in a rodenticide bait matrix that is sufficiently palatable so as to give effective rodenticidal as well as insecticidal activity.

The present invention relates to the systemic insecticide fipronil in conjunction with a rodenticide. It has now been found that high palatability of rodenticide/fipronil baits can be achieved so as to give effective rodenticidal and insecticidal activity as well as acaricidal activity.

Accordingly, in a first aspect, this invention provides use of an effective amount of a rodenticide and an insecticidally effective amount of fipronil in the manufacture of a bait composition for providing a lethal effect on fleas and a host rodent thereof, following ingestion of the bait composition by the host rodent.

In a second aspect, the present invention provides use of an effective amount of a rodenticide and an acaricidally effective amount of fipronil in the manufacture of a bait composition for providing a lethal effect on ticks and a host rodent thereof following ingestion of the bait composition by the host rodent.

For effectiveness against fleas, amounts of fipronil of at least 1 ppm (ppm: parts per million = 0.0001% = 0.001g active/kg of bait) may be present, preferably at least 2 and more preferably at least 10 ppm. For effectiveness against ticks, amounts of fipronil of at least 10 ppm may be present, preferably at least 15 ppm and more preferably at least 25 ppm. The upper limit to the amount of fipronil in the composition will largely be dictated by issues of cost effectiveness and by the risk of inducing toxicosis in the target rodents. Bait with a fipronil concentration of above 200 ppm could cause symptoms of toxicosis in rodents, even in a single feed. It is

not desirable for the fipronil to be present in such amount as it is likely to lead to bait shyness. The term "bait shyness" relates to the scenario where a rodent eats a sub-lethal amount of bait, feels sick and associates its sickness with the bait.

Consequently, the rodent refuses to eat the bait again. Amounts of fipronil up to but not including levels sufficient to cause symptoms of toxicosis are within the scope of the invention. For economic reasons, lower levels may be selected. Amounts of up to 40 ppm of fipronil are suitable against ticks and fleas, being insecticidally and acaricidally effective and generally cost effective.

10 Rodenticides

The rodenticide is a subacute or chronic type rodenticide. These types of rodenticide are slow acting and typically take more than 12 hours, often up to 24 hours or longer, for the onset of symptoms of toxicosis to appear. This is necessary to allow sufficient time for fipronil to be absorbed into the blood of the rodent and for the parasitic arthropods to ingest a lethal dose.

The rodenticide is preferably a hydroxycoumarin or indane-dione anticoagulant. Such anticoagulant rodenticides include, but are not limited to, warfarin, diphacinone, difenacoum, chlorphacinone, flocoumafen, bromadiolone, brodifacoum, difethialone, pindone, coumatetralyl, coumafuryl and coumachlor.

Anticoagulants are suitable rodenticides for preventing bait shyness and for allowing sufficient time for the fipronil to be effective against parasitic arthropods. Anticoagulants minimise the risk of bait shyness because the time between eating bait and feeling sick is in the order of days rather than hours. The time from ingestion of a lethal dose to death is usually in the order of 4-10 days. This is too long a time between eating and the onset of illness for the rodent to make the association.

For the bait composition of the present invention, the rodenticide is preferably selected from the group consisting of brodifacoum, difethialone, flocoumafen, bromadiolone and mixtures thereof.

The rodenticide is preferably included in an amount sufficient to provide a lethal dose if a rodent feeds on the bait for one night only. Amounts providing a lethal dose after a longer period of feeding, however are also within the scope of the

invention. Suitable amounts of brodifacoum, difethialone, flocoumafen and bromadiolone or mixtures thereof are from 10 to 250ppm, preferably 20 to 100 ppm, more preferably 25 to 50ppm.

5 Brodifacoum, bromadiolone, difethialone and flocoumafen are insoluble or weakly soluble in water. Manufacturing concentrates of these anticoagulants traditionally contain the active dissolved in an organic solvent, the solvent being selected to be palatable or at least tasteless to rodents. Or they may be available as a powder pre-mix in which some inert powdered ingredient (eg. wheat flour) is used as a diluent. For the composition of the present invention, it is desirable to maintain
10 neutral pH to avoid hydrolysis of fipronil. The likely effect of any solvent or diluent on pH must therefore be considered. The most preferred rodenticide for the bait composition of the present invention is brodifacoum. Usually, brodifacoum, as its triethanolamine salt, has been provided as a solubilised 0.25% or 2.5% concentrate in a solvent (eg. propylene glycol) with a red or blue dye. Preferably, the brodifacoum
15 for the present invention is formulated without triethanolamine and provided as a 0.25% solution.

Rodenticides which are not anticoagulants but which are also suitable for the present invention include bromethalin, flupropadine, norbormide, calciferol and cholecalciferol (vitamin D3).

20 In a preferred aspect, the present invention provides a rodenticidal bait composition comprising an insecticidally effective amount of fipronil; a rodenticidally effective amount of a rodenticide selected from the group consisting of brodifacoum, difethialone, flocoumafen and mixtures thereof; at least one feeding stimulant and optionally at least one attractant.

25 In a further preferred aspect, the invention provides a rodenticidal bait composition comprising an acaricidally effective amount of fipronil, a rodenticidally effective amount of a rodenticide selected from the group consisting of brodifacoum, difethialone, flocoumafen and mixtures thereof; at least one feeding stimulant and optionally at least one attractant.

30 The bait composition optionally includes one or more insecticidally, acaricidally and rodenticidally compatible excipients and/or adjuvants such as dye, bittering agent, solvent, flow agent, binder, weatherability enhancer and preservative.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

- 5 Combinations of fipronil with brodifacoum have been found to be highly palatable to *Rattus norvegicus* (Norway rat) and *Mus domesticus* (house mouse). It is expected that combinations of fipronil with difethialone, flocoumafen and mixtures thereof including mixtures containing brodifacoum will also be highly palatable to these rodents. It is expected that the combinations will also be palatable to other
- 10 rodents such as *Arvicola terrestris*, *Microtus arvalis*, *Microtus pennsylvanicus*, *Tatera indica*, *Peromyscus leucopus*, *Peromyscus maniculatus*, *Mastomys natalensis*, *Rattus rattus*, *Rattus argentiventer*, *Rattus exulans*, *Sigmodon hispidus*, *Arvicanthis niloticus*, *Bandicota bengalensis*, *Bandicota indica*, *Nesokia indica*, *Meriones hurrianae*, *Millardia meltada* and all members of the *Mus* genera as these rodents are
- 15 also feeders on grain-based materials/baits. Suitable amounts of fipronil and rodenticide are as discussed above.

Attractants/Feeding Stimulants

The bait composition of the present invention includes at least one feeding stimulant and optionally at least one attractant. An attractant is a material that is used to help bring a rodent close to the bait. A feeding stimulant entices the rodent to feed and to keep feeding on the bait. A material may function as both an attractant and a feeding stimulant. Attractants can be a food item or a 'curiosity enhancer'. The former motivates the rodent to approach the bait out of desire to eat it while the latter motivates a rodent to approach out of some other, non-food related desire (eg. to investigate where an interesting odour is coming from). Pheromones (eg. sexual attractants) are examples of curiosity enhancers. Attractants rely on odour solely to bring the rodent into close proximity to the bait thereby increasing the chance that the rodent will eat the bait. Examples of attractants and feeding stimulants that are suitable for the composition of the present invention are given in Table 2 below.

Table 2.

Material	Attractant ?	Feeding Stimulant?
seeds, cereal grains	Yes	Yes
Sugar and sugar products (eg. granulated sugar, confectioner's sugar at 1-5%)	Yes	Yes
Pheromones	Yes	No
Corn starch/corn meal	Yes	Yes
Salt	No	Yes
Monosodium glutamate	No	Yes
Vegetable oils (eg. one or more of peanut, coconut, sesame, sunflower, linseed, palm, rapeseed, olive, corn and soyabean)	Yes	Yes

Oils are also used as 'stickers' in seed/grain bait, binding the poison mix to the outside of the bait.

Suitable stimulants, which are also attractants include: whole and processed seeds including cereal grains; sugar and sugar products; honey; meat and meat products including blood and fat; dairy products; eggs and egg products including shell and yolk; starch; whole and processed nuts; and vegetable oils. The term "seeds" is intended to include seeds in general such as sunflower, thistle, poppy and pumpkin seeds as well as cereal grains such as oats, wheat, rice, barley, corn and millet. The term "processed seeds" is intended to include seeds that are crushed, cracked, rolled, or milled to various consistencies including flour. Suitable vegetable oils include peanut, coconut, sesame, sunflower, linseed, palm, rapeseed, olive, corn, soybean and blends of two or more thereof.

The bait composition of the present invention may include a mixture of feeding stimulants. It may also include at least one attractant. Suitable attractants include pheromones, yeast, and black pepper.

Preferred attractants/stimulants are whole and processed seeds including cereal grains; sugar and sugar products; starch; nuts and processed nuts; and vegetable oil.

Suitably, the total amount of attractants/stimulants in the bait composition as a percentage by weight will range from 50-99.999%.

20 Optional excipients and adjuvants

The bait composition may also comprise one or more optional components such as dye, bittering agent, flow agent, binder, weatherability enhancer and preservative.

Dyes are often added to bait to clearly identify them as non-food items, to deter accidental consumption by people, to deter consumption by non-target animals and to disclose consumption of the bait in the faeces or vomitus. The most common colours used are deep greens and blues.

Bittering agents may be included to minimise the risk of accidental consumption by humans. A suitable agent is denatonium benzoate. This is an extremely bitter tasting compound that at the optimum concentration will be highly distasteful to people but not to rodents. When present, a human taste deterrent is

suitably used in an amount of from 1 to 200 ppm, preferably from 1 to 100 ppm, more preferably from 5 to 50 ppm and most preferably from 5 to 20 ppm.

Flow agents and binders may be included depending on the format of the bait composition. Binders (eg. corn oil) are used to stick the poison to the outside of loose whole grains and seeds or to help provide some cohesion to the bait if prepared in the form of a paste. Flow agents (eg. mineral clay, aluminium silicate) facilitate extrusion and are therefore often used in pellets and extruded blocks. Various bait formats are discussed below.

The addition of paraffin wax to bait greatly improves its resistance to moisture and hence its weatherability. Paraffin greatly improves the effectiveness of rodenticidal baits in tropical, humid climates; in damp indoor locations (kitchens, garages etc.); and in a number of different outdoor situations (eg. sewer and burrow baiting). Suitable amounts of paraffin range from 5-50%.

Other insecticides (eg. 0.1% Malathion) and mould inhibitors (eg. 0.1% 2, 3, 5 trichlorophenylacetate) may be added to grain-based bait to prevent attack by insect pests and to extend the shelf life of the product. Antioxidants may be included to preserve oils and animal products. Antioxidants that may be included are TBHQ, butylated hydroxytoluene or butylated hydroxyanisole. These may be present in amounts of 10 ppm to 20,000 ppm (more ideally 0.05-1%). Amongst the preservatives that may be used are sorbic acid and salts thereof (e.g. potassium sorbate), DowicilTM (Dow-Elanco) and methyl- and propylparabens. Suitably, preservatives may be present in an amount from 0.01-1% more preferably 0.05-0.5%.

Bait Formats

There are a wide variety of conventional bait formats suitable for the present invention. The choice of format will depend on the environment in which it is to be used. No format is ideal for all situations and problems. Table 3 below lists some conventional formats and summarises the advantages and disadvantages of each.

Meal

Meal baits consist of a mixture of whole, ground and/or rolled grains in a range of sizes from fine powdery particles (flour, corn meal) to whole rolled oats, or whole or broken grains.

- 5 Rodents commonly carry food back to their nest or borrow where they may eat it, store it and eat it at a later time, or store it and never eat it. For poisoned bait, the latter is clearly wastage. Meal minimises hoarding simply because it is difficult for a rodent to carry away. The main disadvantages of meal is that it can be messy to use (flour is hard to clean up and can be inhaled), it can be easily contaminated with
- 10 dirt reducing its desirability to rodents, and its quality quickly deteriorates once removed from its packaging.

Format	Advantage	Disadvantage
Meal	<ul style="list-style-type: none"> Highly palatable. Easily digested (small particle size). Hard to hoard (cannot be easily carried away). Economical. 	<ul style="list-style-type: none"> Dust can be easily spread (clean-up is more difficult). Dust can be inhaled by user. Short life-span: affected by moisture (very poor weatherability). Palatability may be an issue if the active is intrinsically unpalatable (poison is on surface). Easily contaminated with dirt which may reduce palatability.
Seed/grain (whole or cut)	<ul style="list-style-type: none"> Highly palatable. Grains can be very economical. 	<ul style="list-style-type: none"> Attractive to non-target, granivorous (grain/seed eating) birds and mammals. Potential poisoning risk to people (children or hungry adults). Palatability may be an issue if the active is intrinsically unpalatable (poison is concentrated on the surface of the seeds/grain and can be wiped off). Poor weatherability. Some seeds may be expensive. Can be easily hoarded.
Pellet	<ul style="list-style-type: none"> More palatable than waxed products. Hard with edges: attractive for rodents to gnaw on. Less attractive to non-target animals (including humans). Low spillage rates – easier clean up. 	<ul style="list-style-type: none"> Can be hoarded. Poor weatherability. Small enough to be taken by granivorous birds.
Waxed pellet	<ul style="list-style-type: none"> Moisture resistant. Good for humid climates or the wetter parts of homes (laundry or garage). Edges for gnawing. Low attractiveness to non-target animals (including people). Low spillage rates – easier clean up. 	<ul style="list-style-type: none"> Probably less palatable than normal pellets. Small enough to be taken by granivorous birds. Can be hoarded.
Moulded wax block	<ul style="list-style-type: none"> Moisture and rain resistant. Can be used outdoors. Multiple gnawing edges. Less attractive to non-target animals (including people). Very low spillage - easy clean up. Smooth, shiny surface is attractive to users. Can be secured to substrate (nailed or tied). 	<ul style="list-style-type: none"> Low palatability due to high wax content. High temperature manufacture: some flavour and freshness is lost and so palatability is reduced. Can melt and soften at high temperatures: palatability is reduced. Smooth, shiny surface is less attractive to rodents.
Compressed or Extruded wax block	<ul style="list-style-type: none"> Moisture resistant. Very low spillage - easy clean up. Low attractiveness to non-target animals (including people). Lower temperatures during manufacturing than wax blocks: better palatability. Rougher surface and more gnawing edges than moulded blocks. Lower wax content than moulded blocks: better palatability. Melting at high temperatures is less of a problem. Can be secured to substrate (nailed or tied). 	<ul style="list-style-type: none"> Low palatability but probably higher than moulded blocks Not as water resistant outdoors as are moulded blocks because less wax is used. Compressed block probably has lower weatherability than extruded block.

Seeds/cereal grains

Seed mixes such as canary seed, and whole, cracked or rolled cereal grains such as wheat, rice, maize, oat, barley and millet form the basis of most commercial baits. For whole grain baits, husks or hulls are removed. The poison is stuck to the outside of the seed or grain using a 'sticker' substance while the inside of the grain or seed may remain free of poison.

Seeds in general and cereal grains in particular are a highly desirable food for commensal rodents. The type of seed or grain preferred depends on the type of rodent and on the type of seed or grain with which they are familiar (rice, wheat, millet etc). The size of the grain or seed should be within preferred range for the target rodent. For example, Norway rats prefer grains in the 0.4 to 0.7 mm diameter size range. Rats generally prefer bigger grains and seeds than mice.

The main disadvantage of whole seed or grain bait is that it is easy for a rodent to hoard and it can be very attractive to nontarget animals such as granivorous (seed or grain eating) birds. Also, this format could have palatability problems if the poison is intrinsically unpalatable. The poison is stuck to the outside of the grain or seed so is at a relatively high concentration. When a rodent bites into the bait it is much more likely to taste a poison than would be the case for more homogenous bait formats such as pellets and blocks. A similar difficulty may apply to meal baits.

Pellets

Pellets are produced by extruding a steamed, soft, hot dough mix of milled grain, poison and other additives through a die after which they are then cut to size.

Pellets are hardened by compression during the process of extrusion and, after extrusion, as they set and moisture content is lowered during oven drying. The degree of hardening is a function of temperature, compression pressure used and drying time. The result is hard, brittle pellets of a consistent diameter and length. Size may vary depending on the target rodent (rat or mouse). Pellets are usually about 3-5 mm diameter and 5-10 mm long cylinders. The addition of an amount of paraffin wax improves the moisture-resistance capabilities (ie. weatherability) of the pellets. However, while this extends their range of applications to, for example, more humid areas of the house and gives the pellets better performance in humid climates,

it also probably lowers the palatability of the bait. Pelletised bait is the most widespread and common rodenticide format and appears to be good general-purpose bait.

5 Moulded Wax Blocks

Moulded blocks are made by pouring a hot blend of grain, melted wax (typically 25-40%), poison and other additives into a mould to produce, upon cooling, a smooth, shiny, solid but waxy product. Blocks of 3-5 g and 15-35 g are typically produced.

- 10 Of the bait formats this is the most moisture resistant but also the least palatable to rodents (the general rule is that the higher the wax content the less palatable the bait). Furthermore, the high temperature used during manufacture cooks and reduces the freshness and palatability of grains in the bait. Exposure to high temperatures (eg. hot climates, or when put in roof voids) can cause the wax to
- 15 soften. This makes the bait unpalatable to rats and mice and so ineffective. The smoothness and shiny appearance of these blocks can make them less attractive to rodents.

- Blocks may be made with a hole through the centre. This allows the blocks to be nailed to a substrate or secured in bait stations to prevent rodents carrying the bait
- 20 away. Holes allow the blocks to be suspended off the ground (eg. nailed to roof rafters in black rat control, or to minimise exposure to water) or threaded onto metal wire for insertion into burrows or down drains (Norway rat control).

Extruded Blocks or Cake

- 25 The palatability problem with moulded blocks is ameliorated with extruded blocks in two ways: less wax is used and the manufacturing temperature is lower. Though less wax is used, extruded blocks still have excellent moisture resistance characteristics. They are produced using a process similar to that used to manufacture bait pellets – extrusion and compression of dough through a die with subsequent
- 30 cutting to size. Compared to moulded blocks, extruded blocks are harder, have a duller, relatively rougher surface; all three features increasing the attractiveness of

these blocks to rodents. Some are designed to be broken into smaller pieces by the user whereas others are cut to a size as they exit the die.

The manufacturing process allows these products to be made into complex shapes with multiple sharp edges to encourage rodents to gnaw the bait. The idea is that rodents like to chew on corners because this gives purchase for their teeth. Dies are therefore used that aim to maximise the number of corners while optimising their arrangement on the block. The effectiveness of blocks with many corners versus those with few is unknown. As for moulded blocks, extruded blocks may be made with hole for attachment. The lower wax content also makes them less prone to softening at high temperatures.

Compressed Blocks

Like extruded blocks, the compressed block format also aims to optimise the balance between palatability and weatherability. They are not made by extrusion through a die but by compression of a warm dough mix in a mould that shapes and compresses it to ensure a hard block upon cooling and drying. Compressed blocks have a dull finish but are smoother than extruded blocks. They also have a dusty surface (perhaps to facilitate their release from the mould).

Paste

This is a soft mix of meal bait based on fats or oils. This format thus has a high moisture content differentiating it from all other formats. It can be a very useful format to use in locations where spillage of bait can be a problem (eg. food storage areas, kitchens etc.) or where rodents have limited access to water. Pastes and gels may be applied with caulking guns.

Comparison of Formats

As mentioned above, each format is generally designed for different rodent control situations and problems. Table 4 compares format on a number of key performance measures.

Table 4. Comparison formats on key performance measures.

Feature	Best format	Worst format
Palatability	Meal or whole seed and grain	Moulded wax block
Moisture resistance	Moulded wax block	Meal
Resistance to hoarding	Meal and block products	Whole seed and grain
Spillage minimisation	Block products	Meal
Economy	Whole grain	Block products/pellets

5

Modes for Carrying Out the Invention

The invention will now be further illustrated with reference to the following non-limiting examples:

10 EXAMPLESPelletised bait compositions

The following example bait formulations use 0.25% w/w or 2.5% w/w brodifacoum liquid concentrates to make a bait with a nominal brodifacoum concentration of 50 ppm (0.005% w/w). Suitable liquid and solid concentrates of other rodenticides (e.g. difethialone at 0.12% w/w liquid concentrate or 0.5% w/w dry concentrate; or flocoumafen at 0.5% dry concentrate) can also be substituted into the formulations. Difethialone is typically formulated as a 25 ppm bait while actives such as flocoumafen and bromadiolone are formulated as 50 ppm baits. If a different active or a concentrate of a particular active is to be used then the baits are made up to 100% by weight by adjusting the percentage of wheat flour added to the formulations.

<u>Component</u>	<u>Wt%</u>	<u>Function</u>
25 <u>Example 1</u>		
Wheat flour	82.892	Food attractant/stimulant

	Millet	10.000	Food attractant/stimulant
	Confectioner's sugar	3.500	Feeding stimulant
	Corn meal	1.500	Food attractant/stimulant
	Brodifacoum 0.25% concentrate	2.000	Rodenticide Active
5	Green dye	0.100	Colour
	Denatonium benzoate (25% solution)	0.004	Bittering agent
	Fipronil	0.004	Insecticide/acaricide

Example 2

10	Wheat flour	88.892	Food attractant/stimulant
	Sugar	5.000	Feeding stimulant
	Corn meal	4.000	Food attractant/stimulant
	Brodifacoum 0.25% concentrate	2.000	Rodenticide Active
15	Green dye	0.100	Colour
	Denatonium benzoate (25% solution)	0.004	Bittering agent
	Fipronil	0.004	Insecticide/acaricide

Example 3

20	Wheat flour	85.392	Food attractant/stimulant
	Corn starch	0.500	Food attractant/stimulant
	Millet	10.000	Food attractant/stimulant
	Confectioner's sugar	2.000	Feeding stimulant
25	Brodifacoum 0.25% concentrate	2.000	Rodenticide Active
	Green dye	0.100	Colour
	Denatonium benzoate (25% solution)	0.004	Bittering agent
	Fipronil	0.004	Insecticide/acaricide

30 Example 4

	Wheat flour	84.392	Food attractant/stimulant
	Millet	5.000	Food attractant/stimulant
	Sugar	4.000	Feeding stimulant
35	Peanut meal	2.500	Food attractant/stimulant
	Corn oil	2.000	Food attractant/stimulant
	Brodifacoum 0.25% concentrate	2.000	Rodenticide Active

Green dye	0.100	Colour
Denatonium benzoate (25% solution)	0.004	Bittering agent
Fipronil	0.004	Insecticide/acaricide

Example 5

5	Wheat flour	87.192	Food attractant/stimulant
	Rolled oats	5.000	Food attractant/stimulant
	Corn meal	5.000	Food attractant/stimulant
	Confectioner's sugar	2.500	Feeding stimulant
10	Brodifacoum 2.5% concentrate	0.200	Rodenticide active
	Green dye	0.100	Colour
	Denatonium benzoate (25% solution)	0.004	Bittering agent
	Fipronil	0.004	Insecticide/acaricide

15 Example 6

	Wheat flour	78.692	Food attractant/stimulant
	Corn Meal	5.000	Food attractant/stimulant
	Millet	10.000	Food attractant/stimulant
20	Confectioner's sugar	4.000	Feeding stimulant
	Peanut Meal	2.000	Food attractant/stimulant
	Brodifacoum 2.5% concentrate	0.200	Rodenticide active
	Green dye	0.100	Colour
	Denatonium benzoate (25% solution)	0.004	Bittering agent
25	Fipronil	0.004	Insecticide/acaricide

- It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered
- 30 in all respects as illustrative and not restrictive.

Method of preparation

The formulations given above was prepared as follows:

- 35 i) Fipronil was weighed into a flask.

- ii) The liquid ingredients were weighed into the flask containing the fipronil. These ingredients were then stirred vigorously for 2 hours on magnetic stirrer or until the fipronil was completely dissolved.
- 5 iii) The remaining dry ingredients were pre-weighed and added to the a powder mixing vessel (Forberg Paddle mixer – 20L capacity).
- iv) The mixer was switched on and the contents allowed to mix until well blended (approximately 5 minutes).
- v) The liquid ingredients were transferred to a dispensing device. This device was specially designed to spray liquid under pressure into the powder mixing
10 vessel.
- vi) After the dry ingredients had mixed for approximately 5 minutes, the liquid ingredients were sprayed into the mixture. The liquid and dry ingredients were allowed to mix for an additional 5 minutes.
- vii) The mixed material (mash) was removed from the powder mixer and poured
15 into the pelletising feed hopper. The pelletiser used was CPM Laboratory Pellet Mill Serial No. 386003
- viii) The mash passes from the feed hopper via a screw conveyer to the conditioner. In this chamber the mash is tumbled in the presence of steam increasing the temperature of the mash to 70-90°C.
- 20 ix) From the conditioner the mash enters the pelletiser where it is compressed between a pressure roller and a rotating die. Continuous rods of compressed mash are extruded from the die to be cut to length by a knife to form the pellets.
- x) The pellets then passed down a chute out of the pelletiser where they are
25 collected and spread onto trays to cool and dry at ambient air temperature for at least 60 minutes.
- xi) The cooled dried pellets were then roughly screened to remove undersized pellets.
- xii) Samples of pellets were taken for laboratory analysis.

Efficacy Results

The following nonlimiting examples provide further demonstration of the utility of the present invention. The bait formulation used in each of these was the same, nominally 50 ppm brodifacoum and 40 ppm fipronil. Different studies did not necessarily use the same batch of material.

Example 7

This study aimed to determine if the invention could kill 90% or more of laboratory strains of Norway rat (*Rattus norvegicus*) (Sprague-Dawley strain) and house mice (*Mus domesticus*) (Swiss-Webster strain) in a single nights exposure. These tests, conduct in June 2003, followed the United States Environmental Protection Agency Office of Pesticide Program Protocols 1.203 (rat) and 1.204 (mouse) modified for a 1-day exposure period. These are choice-feeding trials in which the test animals were simultaneously exposed to the test bait and a non-poisonous but palatable challenge diet. The treatment groups consisted of 20 rats and 20 mice in a 1:1 sex ratio. A control group of 10 rats or 10 mice (also at a 1:1 sex ratio) were also included. The control animals were only exposed to the challenge diet. Following acclimatisation, treatment animals were exposed to the test bait for 24 hours. After 24 h, the test bait was removed and the animals were henceforth fed only on the challenge diet. The animals were monitored daily until any sick animals had either died or recovered. The time of death was recorded for any animal that succumbed during the monitoring period.

No mortality was recorded in any of the control rats or mice. Mortality for rats was 95% with a mean time to death of 7.2 days. Mortality for mice was also 95% with a mean time to death of 6.6 days. Mortality of both rats and mice following a single day exposure to the test bait was $\geq 90\%$ demonstrating that the invention kills laboratory strains of Norway rats and house mice in a single night's feeding.

Example 8

This study determined the efficacy of the invention against nymphal stage ticks *Ixodes trichosuri* on laboratory Norway rats (Wistar Strain). Conducted during June 2003, the trial was a replicated choice-feeding study in which rats had a choice of the invention and a challenge diet over a 3-day exposure period. Nymphal ticks were attached to the rats in a retainer on the shaved neck of the rats 1 day prior to exposure to the bait. The retainers were designed to ensure easy monitoring of ticks with minimal disturbance to both the rats and the ticks. Ticks typically attached within a few hours of introduction onto the rats. The attachment site of each tick was recorded allowing the progress of individual ticks to be monitored. The study consisted of two replicates of 5 male and 5 female rats to each of which a maximum of 8 ticks was attached. Each replicate had a control group of 5 male and 5 female rats to which a maximum of 8 ticks were also attached. The control animals were not exposed to the invention and were fed only on the challenge diet during the monitoring period. The rats and their ticks were monitored daily until any sick rats had either died or recovered. Rats that had become moribund clearly from the effects of the anticoagulant poison were humanely euthanased. The time of death was recorded for each rat and the size and status (alive, dead, moribund) of each ticks was also recorded daily. Any dead ticks were removed and new ticks added when the total number on the rat was 4 or less.

No mortality was recorded for any of the control rats. Tick mortality on control rats was < 5% with ticks needing 4-7 days to engorge and voluntarily detach. The average time to death of the treatment rats was 6.4 days. (This is a slight underestimation of the actual time to death because 65% of the rats were euthanased because they had become terminally moribund. The actual time to death would likely be between 7.0-7.5 days) All the initial cohort of ticks placed on the treatment rats failed to engorge and died within 7 days of exposure to the invention. By the time rats started to die from the effects of brodifacoum (3 days from the first exposure), survival of the initial cohort of ticks was only 41% and was < 5% by the fifth day following first exposure. The death rate for the ticks was therefore faster than that of

the rats. Additional ticks added to surviving rats all failed to engorge and died within 2-3 days of attachment. There was no evidence that ticks failed to attach to treated rats indicating that treatment with the invention did not make the rats repellent to the ticks. These results demonstrate that the invention is effective at killing Norway rats, at killing ticks that infested rats prior to exposure to invention, and at killing ticks that attached to the treated rats after exposure to invention.

10 Example 9

The efficacy of the invention against nymphal paralysis tick *Ixodes holocyclus* on the house mouse (Swiss outbred Strain) was determined in October 2003 using a methodology similar to that described in Example 8. This study, too, was a choice-feeding trial over a 3-day exposure period using ticks held in a retainer on the shaved neck of the mice. Ticks were placed on the mice prior to exposure to the test bait, and fresh ticks introduced as the ticks on the mice died. A single replicate of 5 male and 5 female mice in each of the treatment and control groups was used. A total of 8 ticks were attached to each treatment and control mouse. Additional ticks were added to a mouse whenever the number of surviving ticks was ≤ 4 such that the total number of ticks on the mice at any time did not exceed 8.

Mortality for the control mice reached 100% with the symptoms displayed suggesting that the cause of death was the toxins produced by the ticks. Tick mortality on control mice was $< 5\%$ with ticks engorging in 4-6 days. All treatment mice also died but in this case the symptoms displayed indicate the cause of death being due to anticoagulant toxicosis. The average time to death of the treated mice was longer than for the control mice (6.0 vs 4.8 days). All treated mice had died or had become terminally moribund by 7 days after first exposure to the invention. In contrast to the high survival rate of ticks in the control group, 95% of the initial cohort of ticks placed on the treatment mice died within 6 days of first exposure to the invention, and all failed to engorge. The death rate of the ticks exceeded that for

the mice. Most ticks added to surviving mice after bait exposure died within 3 days of attachment. Only those attaching within 1-2 days of the time of death of the mice managed to survive the death of the host. These results indicate that the invention kills > 90% of mice, kills > 90% of the ticks placed on mice prior to exposure to the invention before the mice died from the effects of the rodenticide and kills > 90% ticks that are attached to mice after exposure to the invention if attachment occurs before 1-2 days of the death of the mouse.

10

Example 10

The efficacy of the invention against cat fleas (*Ctenocephalides felis*) on Norway rats (Wistar Strain) was determined in a choice-feeding trial over a 3-day exposure period. Adult fleas (10 per rat) were held in a retainer on the shaved neck of the rats. Fleas were placed on the rats prior to exposure to the test bait, and fresh fleas introduced when the number of fleas surviving on the rats was ≤ 6 . The maximum number of fleas on a rat at any time did not exceed 10. The study was conducted in two replicated trials in June 2003. Each replicate consisted of a treatment group of 5 male and 5 female rats, and a control group of 5 male and 5 female rats. Both control and treatment animals were infested with fleas. However, treatment animals were exposed to the invention and an alternative challenge diet whereas the control group was only given the challenge diet.

No mortality was recorded among the control rats and the mortality of fleas on the control rats was < 10%. All treatment rats died or had become terminally moribund within 5-7 days after first exposure to the invention. Mortality of the initial cohort of fleas was > 90% within 3 days of first exposure invention. Mortality of the subsequent cohorts of fleas was > 90% within a single day of their introduction and attachment to the rats. This effect continues until the death of the rat. These results show that the invention was highly effective at killing rats and the fleas placed on

them prior to exposure to the invention. The invention was also very effective at killing any fleas placed on, and attaching to rats after exposure to the bait had ceased.

Example 11

5

This study used the same methodology as that described in Example 10 to determine the efficacy of the invention against the stickfast flea (*Echidnophaga gallinacea*) on Norway rats (Wistar Strain). The two replicates of this study were conducted in June 2003.

10

No mortality was recorded among the control rats and the mortality of their fleas was < 10% during the period of the study. All treatment rats died or had become terminally moribund within 4-10 days of first exposure to the invention. Mortality of the initial cohort of fleas was 100% within 3 days of first exposure to the invention.

15

Mortality of the subsequent fleas was > 95% within a day of their introduction and attachment to the rats. This effect continued until the death of the rat. These results show that the invention was highly effective at killing rats and the fleas placed on them prior to exposure to the invention. The invention was also very effective at killing any fleas placed on, and attaching to rats after exposure to the bait had ceased.

20

Example 12

The efficacy of the invention against the cat flea (*Ctenocephalides felis*) on house mice (Swiss outbred Strain) was determined in July 2003 using the same methodology as that described in Example 10 .

25

Three control mice (15%) died during the study, most likely due to the stress or injury during handling. Flea mortality on the control mice was < 5%. All treatment mice died or had become terminally moribund within 5-9 days of first exposure to the invention. Mortality of the fleas placed on the mice prior to exposure to the invention was 100% within 2 days of first exposure. All fleas added thereafter died within a day of their introduction and attachment. This effect continued until the

30

death of the rat. These results show that the invention was highly effective at killing mice and the fleas placed on them prior to exposure to the invention. The invention was also very effective at killing any fleas placed on, and attaching to mice after exposure to the bait had ceased.

5

Example 13

The efficacy of the invention against the stickfast flea (*Echidnophaga gallinacea*) on house mice (Swiss outbred Strain) was determined in June 2003 using the same methodology as that described in Example 12.

10

No control mice died during the study. Flea mortality on the control mice was < 5%. All treatment mice died or had become terminally moribund within 4-9 days of first exposure to the invention. Mortality of the fleas placed on the mice prior to exposure to the invention was 100% within 2 days of first exposure. More than 95% of fleas added thereafter died within a day of their introduction and attachment. This effect continued until the death of the rat. These results show that the invention was highly effective at killing mice and the fleas placed on them prior to exposure to the invention. The invention was also very effective at killing any fleas placed on, and attaching to mice after exposure to the bait had ceased.

15
20

Conclusions

The results of the series of studies conducted show that the invention:

25

- Is effective at killing rats and mice,
- Will kill > 90% of rats and mice following one nights exposure,
- Kills > 90% of the ticks that were infesting rats and mice prior to exposure to the invention.
- Kills > 90% of the ticks that were infesting rats and mice prior to exposure to the invention before the rats or mice die from the effects of the rodenticide.

30

- Kills > 90% of ticks that are placed on rats and mice after exposure to the invention ceases, so long as attachment occurs within 1-2 days of the death of the rodent.
- 5 • Kills > 90% of the fleas that were infesting rats and mice prior to exposure to the invention.
- Kills > 90% of the fleas that were infesting rats and mice prior to exposure to the invention placed before the rodents die from the effects of the rodenticide.
- Kills > 90% of fleas that are that are placed on rats and mice after exposure to the invention. This effect continues until the death of the rodent.

10

Industrial Applicability

The compositions of the present invention may be used to control rodents known to feed on grain-based baits such as *Rattus norvegicus*, *Rattus rattus*, *Rattus argentiventer*, *Rattus exulans*, *Mus sp.* *Arvicola terrestris*, *Microtus arvalis*,
 15 *Microtus pennsylvanicus*, *Tatera indica*, *Peromyscus leucopus*, *Peromyscus maniculatus*, *Mastomys natalensis*, *Sigmodon hispidus*, *Arvicanthis niloticus*, *Bandicota bengalensis*, *Bandicota indica*, *Nesokia indica*, *Meriones hurrinanae*, and *Millardia meltada* and their ectoparasites such as fleas of the type *Ctenocephalides felis*, *Ctenocephalides canis*, *Xenopsylla cheopis*, *Xenopsylla astia*, *Xenopsylla*
 20 *brasiliensis*, *Echidnophaga sp.*, *Pulex irritans*, *Nosopsyllus fasciatus*; and ticks of the type *Dermacentor variabilis*, *Dermacentor andersoni*, *Dermacentor reticulatus*, *Ixodes pacificus*, *Ixodes holocyclus*, *Ixodes ricinus*, *Ixodes persulcatus*, *Ixodes spinipalpis*, *Ixodes scapularis*, *Ixodes hexagonus*, *Amblyomma americanum*, *Rhipicephalus sanguineus*, *R. simus*, *Haemaphysalis leporispalustris*, *Haemaphysalis*
 25 *leachi* and *Ornithodoros sp.*

By killing the parasitic arthropods, successful transfer to a new host following death of a former host is avoided. The risk of infestation of humans and domestic animals is thereby reduced. This may assist in reducing the incidence or risk of contact with the causative agents of such vector-borne diseases as Lyme disease,
 30 plague, Rocky Mountain spotted fever, Colorado tick fever, Kyansanur Forest disease, Kerneroyo, Powassan encephalitis, Russian spring-summer encephalitis,

Crimean-Congo haemorrhagic fever, tick-borne encephalitis, Mediterranean spotted fever, boutonneuse fever, Q fever, North Asian tick typhus, Queensland tick typhus, murine typhus, tick-bite fever, tularaemia, relapsing fever, ehrlichiosis and babesiosis.

CLAIMS:

1. Use of an effective amount of a rodenticide and an insecticidally effective amount of fipronil in the manufacture of a bait composition to provide a lethal effect on fleas and a host rodent thereof, following ingestion of the bait composition by the host rodent.
2. Use according to claim 1 wherein the fipronil is included in the bait composition in an amount of at least 1 ppm.
3. Use according to claim 1 wherein the fipronil is included in the bait composition in an amount of at least 2 ppm.
4. Use according to claim 1 wherein the fipronil is included in the bait composition in an amount of at least 10 ppm.
5. Use according to claim 1 wherein the fipronil is included in the bait composition in an amount of about 40 ppm.
6. Use according to any one of the preceding claims wherein the rodenticide is a hydroxycoumarin or indane-dione anticoagulant.
7. Use according to claim 6 wherein the rodenticide is included in the bait composition in an amount of from 10 to 250 ppm.
8. Use according to claim 6 wherein the rodenticide is included in the bait composition in an amount of from 20 to 100 ppm.
9. Use according to claim 6 wherein the rodenticide is included in the bait composition in an amount of from 25 ppm to 50 ppm.
10. Use according to any one of the preceding claims wherein the rodenticide is selected from the group consisting of brodifacoum, difethialone, flocoumafen, bromadiolone and mixtures thereof.
11. Use according to claim 10 wherein the rodenticide is brodifacoum.
12. Use according to claim 11 wherein the rodenticide is included in the bait composition in an amount of about 50 ppm.
13. Use of an effective amount of a rodenticide and an acaricidally effective amount of fipronil in the manufacture of a bait composition to provide a lethal effect on ticks and a host rodent thereof following ingestion of the bait composition by the host rodent.

14. Use according to claim 13 wherein the fipronil is included in the bait composition in an amount of at least 10 ppm.
15. Use according to claim 13 wherein the fipronil is included in the bait composition in an amount of at least 15 ppm.
- 5 16. Use according to claim 13 wherein the fipronil is included in the bait composition in an amount of at least 25 ppm.
17. Use according to claim 13 wherein the fipronil is included in the bait composition in an amount of about 40 ppm.
18. Use according to any one of claims 13 to 18 wherein the rodenticide is a
10 hydroxycoumarin or indane-dione anticoagulant.
19. Use according to claim 18 wherein the rodenticide is included in the bait composition in an amount of from 10 to 250 ppm.
20. Use according to claim 18 wherein the rodenticide is included in the bait composition in an amount of from 20 to 100 ppm.
- 15 21. Use according to claim 18 wherein the rodenticide is included in the bait composition in an amount of from 25 ppm to 50 ppm.
22. Use according to any one of claims 13 to 22 wherein the rodenticide is selected from the group consisting of brodifacoum, difethialone, flocoumafen, bromadiolone and mixtures thereof.
- 20 23. Use according to claim 22 wherein the rodenticide is brodifacoum.
24. Use according to claim 23 wherein the rodenticide is included in the bait composition in an amount of about 50 ppm.
25. A rodenticidal bait composition comprising:
an insecticidally effective amount of fipronil;
- 25 a rodenticidally effective amount of a rodenticide selected from the group consisting of brodifacoum, difethialone, flocoumafen and mixtures thereof; and
at least one feeding stimulant.
26. A rodenticidal bait composition according to claim 25 further comprising at least one attractant.

27. A rodenticidal bait composition according to claims 25 or 26 further comprising one or more insecticidally and rodenticidally compatible excipients and/or adjuvants selected from the group consisting of dye, bittering agent, solvent, flow agent, binder, weatherability enhancer and preservative.
- 5 28. A rodenticidal bait composition according to any one of claims 25 to 27 comprising, by weight:
at least 0.0001% fipronil;
from 0.001 to 0.025% rodenticide; and
50 to 99.999% of at least one feeding stimulant.
- 10 29. A rodenticidal bait composition according to any one of claims 25 to 28 comprising at least 0.0002% fipronil by weight.
30. A rodenticidal bait composition according to any one of claims 25 to 29 wherein the rodenticide is included in an amount of from 0.002 to 0.010% by weight.
31. A rodenticidal bait composition according to any one of claims 25 to 30
15 wherein the rodenticide is included in an amount of from 0.0025 to 0.0050% by weight.
32. A composition according to any one of claims 25 to 30 wherein the rodenticide is brodifacoum.
33. A composition according to claim 32 wherein the brodifacoum is included in
20 an amount of about 0.0050% by weight.
34. A rodenticidal bait composition comprising:
an acaricidally effective amount of fipronil;
a rodenticidally effective amount of a rodenticide selected from the group consisting of brodifacoum, difethialone, flocoumafen and mixtures thereof; and
25 at least one feeding stimulant.
35. A rodenticidal bait composition according to claim 34 further comprising at least one attractant.
36. A rodenticidal bait composition according to claims 33 or 34 further comprising one or more acaricidally and rodenticidally compatible excipients and/or
30 adjuvants selected from the group consisting of dye, bittering agent, solvent, flow agent, binder, weatherability enhancer and preservative.

37. A rodenticidal bait composition according to any one of claims 34 to 36 comprising, by weight:
at least 0.001% fipronil;
5 from 0.001 to 0.025% rodenticide; and
50 to 99.998% of at least one feeding stimulant.
38. A rodenticidal bait composition according to any one of claims 34 to 37 comprising at least 0.0015% fipronil by weight.
39. A rodenticidal bait composition according to any one of claims 34 to 38
10 comprising at least 0.0025% fipronil by weight.
40. A rodenticidal bait composition according to any one of claims 34 to 39 comprising at least 0.0040% fipronil by weight.
41. A rodenticidal bait composition according to any one of claims 34 to 40 wherein the rodenticide is included in an amount of from 0.002 to 0.010% by weight.
- 15 42. A rodenticidal bait composition according to any one of claims 34 to 41 wherein the rodenticide is included in an amount of from 0.0025 to 0.0050% by weight.
43. A composition according to any one of claims 34 to 42 wherein the rodenticide is brodifacoum.
- 20 44. A composition according to claim 43 wherein the brodifacoum is included in an amount of about 0.0050% by weight.
45. A composition according to any one of claims 25 to 44 wherein the feeding stimulant is one or more items selected from the group consisting of:
whole and processed seeds including cereal grains; sugar and sugar products; honey;
25 meat and meat products including blood and fat; dairy products; eggs and egg products including shell and yolk; starch; whole and processed nuts; and vegetable oils.
46. A composition according to any one of claims 25 to 44 wherein the feeding stimulant is one or more items selected from the group consisting of: whole and
30 processed seeds including cereal grains; sugar and sugar products; starch; whole and processed nuts; and vegetable oil.

47. A rodenticidal bait composition comprising approximately by weight,:
- 0.004 % fipronil:
 - 0.005 % brodifacoum; and
 - 5 50 to 99.7 % of at least one feeding stimulant.
48. A rodenticidal bait composition comprising approximately by weight,:
- 0.004 % fipronil:
 - 0.005 % brodifacoum; and
 - 97 to 99.7 % of at least one feeding stimulant.

ABSTRACTRodenticide

The present invention relates to the systemic insecticide fipronil in conjunction with a rodenticide.

In a first aspect, this invention provides use of an effective amount of a rodenticide and an insecticidally effective amount of fipronil in the manufacture of a bait composition for providing a lethal effect on fleas and a host rodent thereof, following ingestion of the bait composition by the host rodent.

In a second aspect, the invention provides use of an effective amount of a rodenticide and an acaricidally effective amount of fipronil in the manufacture of a bait composition for providing a lethal effect on ticks and a host rodent thereof following ingestion of the bait composition by the host rodent.

In a preferred aspect, the invention provides a rodenticidal bait composition comprising an insecticidally effective amount of fipronil; a rodenticidally effective amount of a rodenticide selected from the group consisting of brodifacoum, difethialone, flocoumafen and mixtures thereof; at least one feeding stimulant and optionally at least one attractant.

In a further preferred aspect, the invention provides a rodenticidal bait composition comprising an acaricidally effective amount of fipronil, a rodenticidally effective amount of a rodenticide selected from the group consisting of brodifacoum, difethialone, flocoumafen and mixtures thereof; at least one feeding stimulant and optionally at least one attractant.